

SERIAL NO: 08/093,972

PATENT

is providing substitute page 38 for printing purposes.

THE SEQUENCE LISTING

A Substitute Sequence Listing and a Declaration have been provided to the examiner, who is requested to substitute it for that filed May 4, 2000.

THE CLAIMS

The claims have been amended as indicated by the examiner and are, therefore, believed to be in allowable form.

Claim 200 has been amended as supported by the specification as filed and the original claims. No new matter is believed to be present in the pending claims. Therefore, the applicant believes they are in condition for allowance.


GENERAL REMARKS

The Assistant Commissioner is hereby authorized to charge \$195.- for a second month extension of time (the first month fee having already been paid), and any other fees owed, or refund any excess, to PTO Deposit Account No. 50-1728, including a fee for an extension of time which, if needed, is hereby requested.

In view of the foregoing amendments and remarks, and of the prior filing of a Substitute Sequence Listing and Declaration, as well as a Petition for Extension of Time and a Notice of Appeal with payment of the requisite fees, this application is believed to be in condition for examination and allowance. Early notice to this respect is solicited.

Respectfully submitted.

EPIGENESIS PHARMACEUTICALS, INC



Viviana Amzel, Ph. D.

Reg. No. 30930

Attorney for the Applicant

August 21, 2001

Date

7 Clarke Drive
Cranbury, NJ 08512
609-409-3035 Ph.
413-845-7530 Fax
Vamzel@epigene.com

s:\legal\00672\Amendment 01-8-21 Supplemental

108. A pharmaceutical composition comprising [,] a carrier; a nucleic acid in the form of an aerosol that comprises one or more oligonucleotide(s) (oligo(s)) effective to alleviate hyper-responsiveness to, and/or increased levels of, adenosine, bronchoconstriction, allergy(ies) and/or inflammation, and contains up to and including about 15% adenosine (A), the oligo being anti-sense to an initiation codon, a coding region or a 5' or 3' intron-exon junctions of a gene encoding an adenosine A₁, A_{2a}, A_{2b} or A₃ receptor or anti-sense to their respective mRNA; pharmaceutically and veterinarily acceptable salts of the oligo(s) or mixtures thereof; and a surfactant that may be operatively linked to the nucleic acid.

109. The composition of claim 108, wherein the oligo consists of up to about 10% A.

110. The composition of claim 109, wherein the oligo consists of up to about 5% A.

111. The composition of claim 110, wherein the oligo consists of up to about 3% A.

112. The composition of claim 111, wherein the oligo is A-free.

113. The composition of claim 108, wherein the oligo is anti-sense to the initiation codon of the mRNA, to the 5' or 3' intron-exon junctions or to sequences of the coding region comprising 2 or more G and/or C of the adenosine A₁ receptor gene.

114. The composition of claim 108, wherein the oligo is anti-sense to the initiation codon of the mRNA, to the 5' or 3' intron-exon junctions or to sequences of the coding region comprising 2 or more G and/or C of the adenosine A_{2a}, A_{2b} and/or A₃ receptors.

115. The composition of claim 108, wherein if the oligo contains adenosine (A), at least one A is substituted by a universal base selected from [the group consisting of] heteroaromatic bases which bind to a thymidine base but have antagonist activity [and] or less than about 0.3 of the adenosine base agonist activity at the adenosine A₁, A_{2b} [and] or A₃ receptors, [and] or heteroaromatic bases [which] that have no activity or have [an] agonist activity at the adenosine A_{2a} receptor.


116. The composition of claim 115, wherein substantially all As are substituted by a universal base (s) selected from heteroaromatic bases that bind to a thymidine base but either have antagonist activity or less than about 0.3 of the adenosine base agonist activity at the adenosine A₁, A_{2b} [and] or A₃ receptors, or heteroaromatic bases that have no activity or have agonist activity at the adenosine A_{2a} receptor.

117. The composition of claim 115, wherein the heteroaromatic bases are selected from pyrimidines or purines that may be substituted by O, halo, NH₂, SH, SO, SO₂, SO₃, COOH, or branched or fused primary or secondary amino, alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, alkoxy, alkenoxy, acyl, cycloacyl, arylacyl, alkynoxy, cycloalkoxy, aroyl, arylthio, arylsulfoxyl, halocycloalkyl, alkylcycloalkyl, alkenylcycloalkyl,

alkynylcycloalkyl, haloaryl, alkylaryl, alkenylaryl, alkynylaryl, arylalkyl, arylalkenyl, arylalkynyl, or arylcycloalkyl, all of which may be further substituted by O, halo, NH₂, primary, secondary or tertiary amine, SH, SO, SO₂, SO₃, cycloalkyl, heterocycloalkyl or heteroaryl.

118. The composition of claim 117, wherein the pyrimidines are substituted at a 1, 2, 3, and/or 4 position, and the purines are substituted at a 1, 2, 3, 4, 7 and/or 8 position.

119. The composition of claim 118, wherein the pyrimidines or purines are selected from theophylline, caffeine, dyphylline, etophylline, acephylline piperazine, bamifylline, enprofylline or xantine having the chemical formula



wherein R¹ and R² are independently H, alkyl, alkenyl or alkynyl and R³ is H, aryl, dicycloalkyl, dicycloalkenyl, dicycloalkynyl, cycloalkyl, cycloalkenyl, cycloalkynyl, O-cycloalkyl, O-cycloalkenyl, O-cycloalkynyl, NH₂-alkylamino-ketoxyalkyloxy-aryl or mono or dialkylaminoalkyl-N-alkylamino-SO₂ aryl.

120. The composition of claim 116, wherein the universal base is selected from 3 - nitropyrrole - 2' - deoxynucleoside, 5 -nitroindole, 2 - deoxyribosyl - (5- nitroindole), 2 - deoxyribofuranosyl - (5 - nitroindole), 2' - deoxyinosine, 2' - deoxynebularine, 6H, 8H - 3, 4 - dihydropyrimido [4, 5 - c] oxazine - 7 - one or 2 - amino - 6 - methoxyaminopurine.

121. The composition of claim 108, wherein a methylated cytosine (mC) is substituted for an unmethylated cytosine (C) in at least one CpG dinucleotide if present in the nucleic acid(s).

122. The composition of claim 108, wherein at least one mononucleotide is linked or modified by one or more of phosphorothioate, phosphorodithioate, phosphorotriothioate, methylphosphonate, phosphoramidate, boranophosphate, phosphotriester, formacetal, 2'-O-methyl, thioformacetal, 5'-thioether, carbonate, 5'-N-carbamate, sulfate, sulfonate, sulfamate, sulfonamide, sulfone, sulfire, sulfoxide, sulfide, hydroxylamine, methylene (methylimino) (MMI) and methyleneoxy (methylimino) (MOMI), terminal 1,3-propanediol, terminal dodecanol, 2'-O-methoxyethyl, C-5-propynyl pyrimidine, C-5 methyl cytidine, C-5 ethynyl pyrimidine, 2'-

SERIAL NO: 08/093,972

PATENT

propoxy, C-18 amine, N3'-P5' phosphoramidates, 3'-alkylamino, 2'-fluoro; 5-fluoro pyrimidine, 5-iodo pyrimidine, 5-bromo pyrimidine, 2'-borano, C-5 hexynyl pyrimidine, 2'-O-(2-methoxy)ethyl, 2'-O-aminopropyl, 5-(phenylethyl) or peptide nucleic acid interbase linkages or conjugated to a polyethylene glycol, cholesterol, cholesteryl, dehydroepiandrosterone (DHEA), dehydroepiandrosterone sulfate (DHEASulfate), dehydroepiandrosterone sulfatide (DHEA Sulfatide), ubiquinone (CoQn), dolichol, poly L-lysine, sulfatidic acid or fatty acids.

123. The composition of claim 122, wherein substantially all mononucleotides are linked or modified by one or more of phosphorothioate, phosphorodithioate, phosphorotrithioate, methylphosphonate, phosphoramidate, boranophosphate, phosphotriester, formacetal, 2'-O-methyl, thioformacetal, 5'-thioether, carbonate, 5'-N-carbamate, sulfate, sulfonate, sulfamate, sulfonamide, sulfone, sulfite, sulfoxide, sulfide, hydroxylamine, methylene (methylimino) (MMI) and methyleneoxy (methylimino) (MOMI), terminal 1,3-propanediol, terminal dodecanol, 2'-O-methoxyethyl, C-5-propynyl pyrimidine, C-5 methyl cytidine, C-5 ethynyl pyrimidine, 2'-propoxy, C-18 amine, N3'-P5' phosphoramidates, 3'-alkylamino, 2'-fluoro; 5-fluoro pyrimidine, 5-iodo pyrimidine, 5-bromo pyrimidine, 2'-borano, C-5 hexynyl pyrimidine, 2'-O-(2-methoxy)ethyl, 2'-O-aminopropyl, 5-(phenylethyl) or peptide nucleic acid interbase linkages or conjugated to a polyethylene glycol, cholesterol, cholesteryl, dehydroepiandrosterone (DHEA), dehydroepiandrosterone sulfate (DHEASulfate), dehydroepiandrosterone sulfatide (DHEA Sulfatide), ubiquinone (CoQn), dolichol, poly L-lysine, sulfatidic acid or fatty acids.

124. The composition of claim 108, wherein the [anti-sense] oligo comprises about 7 to about 60 mononucleotides.

125. The composition of claim 108, wherein the oligo comprises [a sequence selected from] SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5 or SEQ ID NO: 7 to SEQ ID NO: 966, or SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5 or SEQ ID NO: 7 to SEQ ID NO: 966, wherein at least one mononucleotide is linked or modified by one or more of phosphorothioate, phosphorodithioate, phosphorotrithioate, methylphosphonate, phosphoramidate, boranophosphate, phosphotriester, formacetal, 2'-O-methyl, thioformacetal, 5'-thioether, carbonate, 5'-N-carbamate, sulfate, sulfonate, sulfamate, sulfonamide, sulfone, sulfite, sulfoxide, sulfide, hydroxylamine, methylene (methylimino) (MMI) and methyleneoxy (methylimino) (MOMI), terminal 1,3-propanediol, terminal dodecanol, 2'-O-methoxyethyl, C-5-propynyl pyrimidine, C-5 methyl cytidine, C-5 ethynyl pyrimidine, 2'-propoxy, C-18 amine, N3'-P5' phosphoramidates, 3'-alkylamino, 2'-fluoro; 5-fluoro pyrimidine, 5-iodo pyrimidine, 5-bromo pyrimidine, 2'-borano, C-5 hexynyl pyrimidine, 2'-O-(2-methoxy)ethyl, 2'-O-aminopropyl, 5-(phenylethyl) or peptide nucleic acid interbase linkages or conjugated to a polyethylene glycol,

SERIAL NO: 08/093,972

PATENT

cholesterol, cholesteryl, dehydroepiandrosterone (DHEA), dehydroepiandrosterone sulfate (DHEA Sulfate), dehydroepiandrosterone sulfatide (DHEA Sulfatide), ubiquinone (CoQn), dolichol, poly L-lysine, sulfatidic acid or fatty acids.

126. The composition of claim 108, wherein the nucleic acid is linked to an agent that enhances cell internalization or up-take and/or a cell targeting agent.

127. The composition of claim 126, wherein the cell internalization or up take enhancing agent is a transferrin, a asialoglycoprotein or a streptavidin.

128. The composition of claim 126, wherein the cell targeting agent comprises a vector, and the nucleic acid is operatively linked to the vector.

129. The composition of claim 128, wherein the vector comprises a prokaryotic or eukaryotic vector.

130. The composition of claim 108, wherein the surfactant is selected from surfactant protein A, surfactant protein B, surfactant protein C, surfactant protein D, [and] surfactant protein D or active fragments thereof, non-dipalmitoyl disaturated phosphatidylcholine, dipalmitoylphosphatidylcholine, phosphatidylcholine, phosphatidylglycerol, phosphatidylinositol, phosphatidylethanolamine, phosphatidylserine, phosphatidic acid, ubiquinones, lysophosphatidylethanolamine, lysophosphatidylcholine, palmitoyl-lysophosphatidylcholine, dehydroepiandrosterone, dolichols, sulfatidic acid, glycerol-3-phosphate, dihydroxyacetone phosphate, glycerol, glycerol-3-phosphocholine, dihydroxyacetone, palmitate, cytidine diphosphate (CDP) diacylglycerol, CDP choline, choline, choline phosphate, lamellar bodies, omega-3 fatty acids, polyenic acid, polyenoic acid, lecithin, palmitic acid, non-ionic ethylene and/or propylene oxide block copolymers, polyoxypropylene, polyoxyethylene, poly (vinyl amine) with dextran and/or alkanoyl side chains, polyoxy ethylene ethers, phenoxy polyethoxy alcohols, phosphatidyl choline esters, [and] phosphatidyl ethers, palmitates, [alcohols] tyloxapol, phospholipids, [neutral lipids,] fatty acids, [or] surfactant-associated proteins or $C_{22}H_{19}C_{10}$.

131. The composition of claim 130, wherein the surfactant is selected from polyoxy ethylene 23 lauryl ether (Brij 35[®]), t-octyl phenoxy polyethoxy ethanol (Triton X-100[®]), dipalmitoyl phosphatidyl choline (DPPC) and phosphatidyl glycerol (PG) (ALEC[®]), tyloxapol (Exosurf[®]), phospholipids, fatty acids, surfactant-associated proteins (Survanta[®]) or $C_{22}H_{19}C_{10}$ (Atovaquone[®]).

133. The composition of claim 108, wherein the carrier comprises a biologically acceptable carrier.

134. The composition of claim 108, wherein the carrier is a pharmaceutically or

SERIAL NO: 08/093,972

PATENT

veterinarily acceptable carrier.

135. The composition of claim 134, wherein the carrier is selected from gaseous, liquid or solid carriers.

136. The composition of claim 108, further comprising an agent selected from therapeutic agents other than the nucleic acid(s), antioxidants, flavoring or coloring agents, fillers, volatile oils, buffering agents, dispersants, RNA inactivating agents, flavoring agents, propellants or preservatives.

137. The composition of claim 136, comprising a pharmaceutically or veterinarily acceptable carrier, the nucleic acid, a surfactant, and other therapeutic agents [a therapeutic agent selected from adenosine A1, A2b or A3 receptor activity inhibiting agents other than the oligo(s), anti-arrhythmic agents, anti-inflammatory agents, anti-bacterial agents, anti-sepsis agents, adenosine or agents exhibiting adenosine agonist activity, analgesics, diuretics, kidney activity maintenance or restoration agents or agents for the treatment of pulmonary vasoconstriction, inflammation, allergies, asthma, acute respiratory distress syndrome (ARDS), ischemia, impeded and blocked respiration, respiratory distress syndrome (RDS), pain, cystic fibrosis (CF), pulmonary hypertension, pulmonary vasoconstriction, emphysema, chronic obstructive pulmonary disease (COPD)].

138. The composition of claim 136, wherein the RNA inactivating agent comprises an enzyme.

139. The composition of claim 138, wherein the enzyme comprises a ribozyme.

140. The composition of claim 108, further comprising a propellant.

141. The composition of claim 108, wherein the nucleic acid is present in an amount of about 0.01 to about 99.99 w/w of the composition.

143. The formulation of claim 108, selected from intrabuccal, intrapulmonary, respirable, nasal, inhalable, intracavitary, intraorgan, or slow release formulations.

144. The formulation of claim 143, wherein the carrier is selected from a gaseous, solid or liquid carrier.

146. The aerosol or spray formulation of claim 108, [wherein] which is selected from powders, sprays, solutions, suspensions or emulsions.

148. The aerosol or spray formulation of claim 108, which is selected from aqueous or alcoholic solutions or suspensions, oily solutions or suspensions, or oil-in-water or water-in-oil emulsions.

151. A capsule or cartridge, comprising the formulation of claim 143.

152. The spray or aerosol formulation of claim 146, comprising a solid powdered spray or aerosol.

153. The formulation of claim 108, wherein the carrier comprises a hydrophobic carrier.

154. The formulation of claim 153, wherein the carrier comprises lipid vesicles and/or particles.

155. The formulation of claim 154, wherein the vesicles comprise liposomes, and the particles comprise microcrystals.

156. The formulation of claim 155, wherein the vesicles comprise liposomes that comprise the nucleic acid.

158. The formulation of claim 143, which comprises an intrapulmonary, intracavitary or intraorgan liquid or solid powdered formulation of particle size about 0.5 μ to about 10 μ , or about 10 μ to about 500 μ .

159. The formulation of claim 143, which comprises a nasal formulation of particle size about 10 μ to about 500 μ .

161. The formulation of claim 143, in bulk, or in single or multiple unit dose form.

162. The formulation of claim 143, which is a respirable or inhalable formulation [of] comprising a solid powdered or liquid aerosol or spray of particle size about 0.5 μ to about 10 μ .

163. A single cell, comprising the nucleic acid of claim 108.

164. A kit for diagnosis or treatment of diseases and conditions associated with hypersensitivity to and/or increased levels of, adenosine and/or bronchoconstriction and/or lung allergy(ies) and/or inflammation and/or asthma comprising, in separate containers, the delivery device of claim 222;

a nucleic acid comprising at least one oligonucleotide (oligo) effective to alleviate hyper-responsiveness to, and/or increased levels of, adenosine, or to alleviate bronchoconstriction, asthma or lung allergy(ies) and/or inflammation, the oligo being anti-sense to the initiation codon, the coding region or the 5' or 3' intron-exon junctions of a gene encoding a protein associated with hyper-responsiveness to, and/or increased levels of, adenosine, with bronchoconstriction, asthma, or lung allergy(ies) or inflammation, or being anti-sense to the corresponding mRNA; the nucleic acid comprising one or more oligo(s), their mixtures or their pharmaceutically or veterinarily acceptable salts [of the oligo(s)]; and

instructions for preparation of a respirable, inhalable, nasal, intrapulmonary, intraorgan, or intracavitary formulation of particle size about 0.5 to about 500 μ and for its use; and

SERIAL NO: 08/093,972

PATENT

optionally an agent selected from therapeutic or diagnostic agents other than the oligo, anti-oxidants, fillers, volatile oils, dispersants, anti-oxidants, flavoring agents, propellants, preservatives, solvents, buffering agents, RNA inactivating agents, agents that are internalized or up-taken by a cell, or coloring agents.

165. The kit of claim 164, wherein the delivery device comprises a nebulizer that delivers single metered doses of a solid powdered or liquid aerosol or spray formulation of particle size about 0.5 μ to about 10 μ or about 10 μ to about 500 μ of the nucleic acid.

166. The kit of claim 164, wherein the device comprises an insufflator adapted for receiving and piercing or opening a capsule(s) or cartridge(s) and producing a solid powdered or liquid aerosol or spray; and the nucleic acid is provided separately in a pierceable or openable capsule(s) or cartridge(s) as a nasal, inhalable, respirable, intrapulmonary, intracavitary or intraorgan formulation of particle size about 0.5 μ to about 10 μ or about 10 μ to about 500 μ .

167. The kit of claim 164, wherein the delivery device comprises a pressurized [inhalator] inhaler that delivers a solid powdered or liquid aerosol or spray of particle size about 0.5 μ to about 10 μ or about 10 μ to about 500 μ ; and the nucleic acid is provided as a suspension, solution, emulsion or dry powder aerosol or spray formulation of about 0.5 μ to about 10 μ or about 10 μ to about 500 μ .

168. The kit of claim 164, comprising the delivery device, a surfactant, the nucleic acid and other therapeutic agents [a therapeutic agent selected from adenosine A1, A2b or A3 receptor antagonists other than the oligo(s), adenosine A2a receptor stimulants, anti-inflammatory agents, anti-histaminic agents, anti-allergic agents, anti-bacterial, anti-virals, analgesics, kidney activity maintenance or restoration agents, anti-cancer agents, adenosine, blood pressure controlling agents, or diuretics].

169. The kit of claim 164, wherein the solvent is selected from organic solvents or organic solvents mixed with one or more co-solvents.

170. The kit of claim 164, wherein the device is adapted for receiving a capsule(s) or cartridge(s), and the nucleic acid is separately provided as an inhalable, respirable, nasal, intracavitary, intraorgan or intrapulmonary formulation in a capsule(s) or cartridge(s).

171. The kit of claim 164 further comprising, in [a] separate containers, a propellant, and pressurized means for delivery adapted for delivering a solid powdered or liquid aerosol or spray, and instructions for loading into the delivery device [the nucleic acid as] an inhalable, respirable, nasal, intracavitary, intraorgan or intrapulmonary formulation of the nucleic acid of particle size about 0.5 μ to about 10 μ or about 10 μ to about 500 μ , and then joining the device with the propellant and the pressurized means.

SERIAL NO: 08/093,972

PATENT

172. The kit of claim 167, wherein the pressurized [inhalator] inhaler further comprises a propellant and means for delivery of the propellant, and delivers the nucleic acid as a liquid or solid powdered aerosol or spray formulation [of the nucleic acid].

173. An in vivo method of delivering a pharmaceutical composition to a target polynucleotide, comprising administering to the airways of a subject an aerosol or spray composition of particle size about 0.5 μ to about 500 μ [,] comprising a nucleic acid which comprises at least one oligonucleotide (oligo) effective to alleviate hyper-responsiveness to, and/or increased levels of, adenosine, or to alleviate bronchoconstriction, asthma and/or lung allergy(ies) and/or inflammation, the oligo containing up to and including about 15% adenosine (A), and being anti-sense to the initiation codon, the coding region or the 5' or 3' intron-exon junctions of a gene encoding a protein associated with hyper-responsiveness to, and/or increased levels of, adenosine, [with] bronchoconstriction, asthma [,] and/or lung allergy(ies) and/or inflammation, or being anti-sense to the corresponding mRNA; the nucleic acid comprising one or more oligo(s), pharmaceutically [and] or veterinarily acceptable salts of the oligo(s), or mixtures of the oligo(s) or their salts.

178. The method of claim 173, wherein the composition is administered intrapulmonarily, intraorgan, intracavitarily, intrabuccally, intranasally, by inhalation or into the subject's respiratory system.

179. The method of claim 173, wherein the oligo is effective to reduce hyper-responsiveness to adenosine, the amount of the adenosine receptor or the production or availability of adenosine, or to increase the degradation of the adenosine receptor mRNA.

180. The method of claim 178, wherein the oligo is administered directly into the subject's lung (s), intraorgan, intracavitarily, intrabuccal or intrapulmonarily.

181. The method of claim 178, wherein the composition is administered as [powdered] solid powdered or liquid particles of the nucleic acid about 0.5 to about 10 μ in size.

184. The method of claim 173, wherein the composition further comprises a surfactant.

185. The method of claim 173, wherein the hyper-responsiveness to, and/or increased levels of, adenosine, [or bronchoconstriction] asthma or lung allergy(ies) or inflammation is associated with bronchoconstriction of lung airways.

186. The method of claim 185, wherein the hyper-responsiveness to, or increased levels of, adenosine, [or] bronchoconstriction, asthma or lung allergy(ies) or inflammation is associated with COPD, asthma, ARDS, RDS, CF or side effects of adenosine administration.

187. The method of claim 173, wherein the hyper-responsiveness to, or increased

SERIAL NO: 08/093,972

PATENT

levels of, adenosine, [or] bronchoconstriction, asthma, or lung allergy(ies) or inflammation is associated with inflammation or an inflammatory disease.

188. The method of claim 173, wherein the composition further comprises other therapeutic agents

[a therapeutic agent selected from adenosine A₁, A_{2b} or A₃ receptor inhibiting agents or adenosine A_{2a} receptor stimulating agents other than the nucleic acid(s), anti-inflammatory agents, anti-bacterial agents, anti-sepsis agents, kidney activity maintenance or restoration agents or agents for the treatment of pulmonary vasoconstriction, inflammation, allergies, asthma, impeded respiration, respiratory distress syndrome (RDS), acute respiratory distress syndrome (ARDS), pain, cystic fibrosis (CF), pulmonary hypertension, pulmonary vasoconstriction, emphysema, or chronic obstructive pulmonary disease (COPD)].

189. The method of claim 188, wherein the therapeutic agent [is selected from] comprises anti-adenosine A₁, A_{2b} or A₃ receptor agents or adenosine A_{2a} receptor stimulating agents other than the nucleic acid(s).

191. The method of claim 184, wherein the surfactant comprises surfactant protein A, surfactant protein B, surfactant protein C, surfactant protein D or active fragments thereof, non-dipalmitoyl disaturated phosphatidylcholine, dipalmitoylphosphatidylcholine, phosphatidylcholine, phosphatidylglycerol, phosphatidylinositol, phosphatidylethanolamine, phosphatidylserine, phosphatidic acid, ubiquinones, lysophosphatidylethanolamine, lysophosphatidylcholine, palmitoyl-lysophosphatidylcholin, dehydroepiandrosterone, dolichols, sulfatidic acid, glycerol-3-phosphate, dihydroxyacetone phosphate, glycerol, glycerol-3-phosphocholine, dihydroxyacetone, palmitate, cytidine diphosphate (CDP) diacylglycerol, CDP choline, choline, choline phosphate, lamellar bodies, omega-3 fatty acids, polyenic acid, polyenoic acid, lecithin, palmitic acid, non-ionic ethylene and/or propylene oxide block copolymers, polyoxypropylene, polyoxyethylene, poly (vinyl amine) with dextran and/or alkanoyl side chains, polyoxy ethylene ethers, phenoxy polyethoxy alcohols, phosphatidyl choline esters, [and] phosphatidyl ethers, palmitates, alcohols and tyloxapol, phospholipids, fatty acids, [or] surfactant-associated proteins, [and] or C₂₂H₁₉C₁₀.

192. The method of claim 173, wherein the subject is a mammal.

193. The method of claim 192, wherein the mammal is a human or a non-human mammal.

195. The method of claim 173, wherein the nucleic acid is administered in amount of about 0.005 to about 150 mg/kg body weight.

196. The method of claim 195, wherein the nucleic acid is administered in an amount

SERIAL NO: 08/093,972

PATENT

of about 0.01 to about 75 mg/kg body weight.

197. The method of claim 196, wherein the nucleic acid is administered in an amount of about 1 to about 50 mg/kg body weight.

198. The method of claim 173, which is a prophylactic or therapeutic method.

200. The method of claim 173, wherein the nucleic acid is obtained by

(a) selecting fragments of a target nucleic acid having at least 4 contiguous bases consisting of G or C; and

[(b) obtaining a first oligo 4 to 60 nucleotides long which comprises the selected fragment and has a C and G nucleic acid content of up to and including about 15%; and (c)]

(b) obtaining a second oligo 4 to 60 nucleotides long comprising a sequence [which] that is anti-sense to the selected fragment, the second oligo having an A base content of up to and including about 15%.

201. The method of claim 173, wherein the oligo consists of up to about 10% A.

202. The method of claim 201, wherein the oligo consists of up to about 5% A.

203. The method of claim 201, wherein the oligo consists of up to about 3% A.

204. The method of claim 203, wherein the oligo is A-free.

205. The method of claim 173, wherein the oligo is anti-sense to the initiation codon, the coding region or the 5' or 3' intron-exon junctions of a gene encoding an adenosine A₁, A_{2b} or A₃ receptor, and the composition further comprises a surfactant.

206. The method of claim 173, wherein if the oligo contains A, at least one A is substituted with a universal base selected from heteroaromatic bases which bind to a thymidine base but have antagonist activity or less than about 0.3 of the adenosine base agonist activity at the adenosine A₁, A_{2b} or A₃ receptors, or heteroaromatic bases which have no activity or have [an] agonist activity at the adenosine A_{2a} receptor.

207. The method of claim 206, wherein substantially all As are substituted with universal bases selected from heteroaromatic bases which bind to a thymidine base but have antagonist activity or less than about 0.3 of the adenosine base agonist activity at the adenosine A₁, A_{2b} or A₃ receptors, heteroaromatic bases which have no activity or have an agonist activity at the adenosine A_{2a} receptor.

208. The method of claim 206, wherein the heteroaromatic bases are selected from pyrimidines [and] or purines [, which] that may be substituted by O, halo, NH₂, SH, SO, SO₂, SO₃, COOH branched fused primary secondary amino, alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, alkoxy, alkenoxy, acyl, cycloacyl, arylacyl, alkynoxy, cycloalkoxy, aroyl, arylthio, arylsulfoxyl, halocycloalkyl, alkylcycloalkyl, alkenylcycloalkyl,


SERIAL NO: 08/093,972

PATENT

alkynylcycloalkyl, haloaryl, alkylaryl, alkenylaryl, alkynylaryl, arylalkyl, arylalkenyl, arylalkynyl, arylcycloalkyl, all of which may be further substituted by O, halo, NH₂, primary, secondary and tertiary amine, SH, SO, SO₂, SO₃, cycloalkyl, heterocycloalkyl or heteroaryl.

209. The method of claim 208, wherein the pyrimidines are substituted at positions 1, 2, 3 and/or 4, and the purines are substituted at positions 1, 2, 3, 4, 7 and/or 8.

210. The method of claim 209, wherein the pyrimidines and purines are selected from theophylline, caffeine, dyphylline, etophylline, acephylline piperazine, bamifylline, enprofylline or xantine having the chemical formula



wherein R¹ and R² are independently H, alkyl, alkenyl or alkynyl, and R³ is H, aryl, dicycloalkyl, dicycloalkenyl, dicycloalkynyl, cycloalkyl, cycloalkenyl, cycloalkynyl, O-cycloalkyl, O-cycloalkenyl, O-cycloalkynyl, NH₂-alkylamino-ketoxyalkyloxy-aryl or mono or dialkylaminoalkyl-N-alkylamino-SO₂ aryl.

211. The method of claim 206, wherein the universal base [is selected from] comprises 3-nitropyrrole-2'-deoxynucleoside, 5-nitro-indole, 2-deoxyribosyl-(5-nitroindole), 2-deoxyribofuranosyl-(5-nitroindole), 2'-deoxyinosine, 2'-deoxynebularine, 6H, 8H-3,4-dihydropyrimido [4,5-c] oxazine-7-one or 2-amino-6-methoxyaminopurine.

212. The method of claim 173, further comprising methylating at least one cytosine vicinal to a guanosine into a methylated cytosine (^mC) if a CpG dinucleotide [if] is present in the oligo(s).

213. The method of claim 173, further comprising modifying or substituting at least one mononucleotide of the anti-sense oligo(s) with methylphosphonate, phosphotriester, phosphorothioate, phosphorodithioate, boranophosphate, formacetal, thioformacetal, thioether, carbonate, carbamate, sulfate, sulfonate, sulfamate, sulfonamide, sulfone, sulfite, sulfoxide, sulfide, hydroxylamine, methylene(methylimino), methyleneoxy (methylimino), 2'-O-methyl, phosphoramidate residues, or combinations thereof.

214. The method of claim 213, wherein substantially all mononucleotides are

SERIAL NO: 08/093,972

PATENT

substituted and/or modified.

215. The method of claim 173, further comprising operatively linking the nucleic acid to an agent [selected from agents] that enhances cell internalization or up-take, or a cell targeting agent[s].

216. The method of claim 215, wherein the cell internalization or up-take enhancing agent is selected from transferrin, asialoglycoprotein or streptavidin.

217. The method of claim 215, wherein the cell targeting agent comprises a vector.

218. The method of claim 217, wherein the vector to which the agent is operatively linked comprises a prokaryotic or eukaryotic vector.

219. The method of claim 173, wherein the nucleic acid comprises an oligo [sequence selected from] of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5 or SEQ ID NO: 7 to SEQ ID NO: 966, or SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5 or SEQ ID NO: 7 to SEQ ID NO: 966, wherein at least one mononucleotide is linked or modified by one or more of phosphorothioate, phosphorodithioate, phosphorotriothioate, methylphosphonate, phosphoramidate, boranophosphate, phosphotriester, formacetal, 2'-O-methyl, thioformacetal, 5'-thioether, carbonate, 5'-N-carbamate, sulfate, sulfonate, sulfamate, sulfonamide, sulfone, sulfite, sulfoxide, sulfide, hydroxylamine, methylene (methylimino) (MMI) and methyleneoxy (methylimino) (MOMI), terminal 1,3-propanediol, terminal dodecanol, 2'-O-methoxyethyl, C-5-propynyl pyrimidine, C-5 methyl cytidine, C-5 ethynyl pyrimidine, 2'-propoxy, C-18 amine, N3'-P5' phosphoramidates, 3'-alkylamino, 2'-fluoro; 5-fluoro pyrimidine, 5-iodo pyrimidine, 5-bromo pyrimidine, 2'-borano, C-5 hexynyl pyrimidine, 2'-O-(2-methoxy)ethyl, 2'-O-aminopropyl, 5-(phenylethyl) or peptide nucleic acid interbase linkages or conjugated to a polyethylene glycol, cholesterol, cholesteryl, dehydroepiandrosterone (DHEA), dehydroepiandrosterone sulfate (DHEASulfate), dehydroepiandrosterone sulfamide (DHEA Sulfamide), ubiquinone (CoQn), dolichol, poly L-lysine, sulfatidic acid or fatty acids.

220. The method of claim 191, wherein the surfactant is selected from polyoxy ethylene 23 lauryl ether (Brij 35[®]), t-octyl phenoxy polyethoxy ethanol (Triton X-100[®]), dipalmitoyl phosphatidyl choline (DPPC) and phosphatidyl glycerol (PG) (ALEC[®]), tyloxapol (Exosurf[®]), phospholipids, fatty acids, surfactant-associated proteins (Survanta[®]) or C₂₂H₁₉C₁₀ (Atovaquone[®]).

221. The method of claim 173, wherein the hyper-responsiveness to, or increased levels of, adenosine, [or] bronchoconstriction, or lung allergy(ies) or inflammation, is associated with asthma or a disease or condition associated with asthma.

222. A diagnostic or therapeutic device adapted for delivering a respirable, inhalable,

SERIAL NO: 08/093,972

PATENT

222. A diagnostic or therapeutic device adapted for delivering a respirable, inhalable, nasal, intrapulmonary, intraorgan, or intracavitary formulation of particle size about $0.5\ \mu$ to about $500\ \mu$, the formulation comprising a nucleic acid [which] that comprises at least one oligonucleotide (oligo) effective for diagnosing or treating hyper-responsiveness to, or increased levels of, adenosines, [or] bronchoconstriction, asthma, or lung allergy(ies) or inflammation, or a disease or condition associated with either of them, the oligo being anti-sense to the initiation codon, the coding region or the 5' or 3' intron-exon junctions of a gene encoding a protein associated with hyper-responsiveness to, or increased levels of, adenosine, bronchoconstriction, asthma, or lung allergy(ies) or inflammation, or being anti-sense to the corresponding mRNA; the nucleic acid comprising one or more oligo(s), their mixtures, or their pharmaceutically or veterinarily acceptable salts.

223. The device of claim 222, comprising a nebulizer adapted for delivering single metered doses of the formulation as a solid powdered or liquid aerosol or spray of particle size about $0.5\ \mu$ to about $10\ \mu$ or about $10\ \mu$ to about $500\ \mu$.

224. The device of claim 222, which comprises an insufflator adapted for receiving and piercing or opening a capsule(s) or cartridge(s) and for producing a solid powdered or liquid aerosol or spray of particle size about $0.5\ \mu$ to about $10\ \mu$ or about $10\ \mu$ to about $500\ \mu$, and wherein the formulation is provided separately in a piercable or openable capsule(s) or cartridge(s) as a nasal, inhalable, respirable, intrapulmonary, intracavitary or intraorgan formulation of particle size about $0.5\ \mu$ to about $10\ \mu$ or about $10\ \mu$ to about $500\ \mu$.

225. The device of claim 222, which comprises a pressurized [inhalator] inhaler that delivers a solid powdered or liquid aerosol or spray formulation of particle size about $0.5\ \mu$ to about $10\ \mu$ or about $10\ \mu$ to about $500\ \mu$; [and] wherein the formulation comprises a suspension, solution, emulsion or dry powder aerosol or spray formulation of the nucleic acid of particle size about $0.05\ \mu$ to about $50\ \mu$ or about $10\ \mu$ to about $500\ \mu$.

226. The pressurized [inhalator] inhaler of claim 225 further comprising, in [a] separate containers, a propellant and pressurized means for delivery[,] adapted for delivering a solid powdered or liquid aerosol or spray, and instructions for loading into [the] the delivery device the inhalable, respirable, nasal, intracavitary, intraorgan or intrapulmonary formulation, and joining the device with the propellant and the pressurized delivery means.

227. The pressurized [inhalator] inhaler of claim 225, further comprising a propellant and propellant delivery means, wherein the pressurized [inhalator] inhaler delivers the formulation as a liquid or solid powdered aerosol or spray.

228. The device of claim 222, which is adapted for receiving and piercing or opening a

SERIAL NO: 08/093,972

PATENT

capsule(s) or cartridge(s), and wherein the formulation is provided separately in a capsule(s) or cartridge(s).

229. The kit of claim 164, wherein the oligo is anti-sense to the initiation codon, the coding region or the 5' or 3' region of a gene encoding a polypeptide selected from an adenosine A₁ receptor, adenosine A_{2a} receptor, adenosine A_{2b} receptor, or adenosine A₃ receptor.

230. The kit of claim 229, for diagnosis or treatment of sepsis, pulmonary vasoconstriction, inflammation, allergies, asthma, impeded respiration, respiratory distress syndrome (RDS), acute respiratory distress syndrome (ARDS), pain, cystic fibrosis (CF), pulmonary hypertension, pulmonary vasoconstriction, emphysema or chronic obstructive pulmonary disease (COPD).

231. The kit of claim 164, wherein the nucleic acid comprises an oligo [sequence selected from] of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5 or SEQ ID NO: 7 to SEQ ID NO: 996, or SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5 or SEQ ID NO: 7 to SEQ ID NO: 996, wherein at least one mononucleotide is linked or modified by one or more of phosphorothioate, phosphorodithioate, phosphorotriothioate, methylphosphonate, phosphoramidate, boranophosphate, phosphotriester, formacetal, 2'-O-methyl, thioformacetal, 5'-thioether, carbonate, 5'-N-carbamate, sulfate, sulfonate, sulfamate, sulfonamide, sulfone, sulfite, sulfoxide, sulfide, hydroxylamine, methylene (methylimino) (MMI) and methyleneoxy (methylimino) (MOMI), terminal 1,3-propanediol, terminal dodecanol, 2'-O-methoxyethyl, C-5-propynyl pyrimidine, C-5 methyl cytidine, C-5 ethynyl pyrimidine, 2'-propoxy, C-18 amine, N3'-P5' phosphoramidates, 3'-alkylamino, 2'-fluoro; 5-fluoro pyrimidine, 5-iodo pyrimidine, 5-bromo pyrimidine, 2'-borano, C-5 hexynyl pyrimidine, 2'-O-(2-methoxy)ethyl, 2'-O-aminopropyl, 5-(phenylethyl) or peptide nucleic acid interbase linkages or conjugated to a polyethylene glycol, cholesterol, cholesteryl, dehydroepiandrosterone (DHEA), dehydroepiandrosterone sulfate (DHEA Sulfate), dehydroepiandrosterone sulfatide (DHEA Sulfatide), ubiquinone (CoQn), dolichol, poly L-lysine, sulfatidic acid or fatty acids.

232. The composition of claim 108, which comprises particle sizes of about 0.5 μ to about 10 μ or about 10 μ to about 500 μ .

233. The nucleic acid of claim 108, which is operatively linked to a vector.

234. A single cell, comprising the nucleic acid of claim 233.